

Dyson-Schwinger Equations – Achievements and Challenges

Craig D. Roberts

cdroberts@anl.gov

Physics Division

Argonne National Laboratory

Dyson-Schwinger Equations – Achievements and Challenges

Peking University, Beijing, China, 13-17 August, 2007

p. 1/46


Study Structure via Nucleon Form Factors



Study Structure via Nucleon Form Factors

- Electron's relativistic electromagnetic current:

$$\begin{aligned} j_\mu(P', P) &= ie \bar{u}_e(P') \Lambda_\mu(Q, P) u_e(P), \quad Q = P' - P \\ &= ie \bar{u}_e(P') \gamma_\mu(-1) u_e(P) \end{aligned}$$




Study Structure via Nucleon Form Factors

- Electron's relativistic electromagnetic current:

$$\begin{aligned} j_\mu(P', P) &= ie \bar{u}_e(P') \Lambda_\mu(Q, P) u_e(P), \quad Q = P' - P \\ &= ie \bar{u}_e(P') \gamma_\mu(-1) u_e(P) \end{aligned}$$

- Nucleon's relativistic electromagnetic current:



Study Structure via Nucleon Form Factors

- Electron's relativistic electromagnetic current:

$$\begin{aligned} j_\mu(P', P) &= ie \bar{u}_e(P') \Lambda_\mu(Q, P) u_e(P), \quad Q = P' - P \\ &= ie \bar{u}_e(P') \gamma_\mu(-1) u_e(P) \end{aligned}$$

- Nucleon's relativistic electromagnetic current:

$$\begin{aligned} J_\mu(P', P) &= ie \bar{u}_p(P') \Lambda_\mu(Q, P) u_p(P), \quad Q = P' - P \\ &= ie \bar{u}_p(P') \left(\gamma_\mu F_1(Q^2) + \frac{1}{2M} \sigma_{\mu\nu} Q_\nu F_2(Q^2) \right) u_p(P) \end{aligned}$$

$$G_E(Q^2) = F_1(Q^2) - \frac{Q^2}{4M^2} F_2(Q^2), \quad G_M(Q^2) = F_1(Q^2) + F_2(Q^2).$$



Study Structure via Nucleon Form Factors

- Electron's relativistic electromagnetic current:

$$\begin{aligned} j_\mu(P', P) &= ie \bar{u}_e(P') \Lambda_\mu(Q, P) u_e(P), \quad Q = P' - P \\ &= ie \bar{u}_e(P') \gamma_\mu(-1) u_e(P) \end{aligned}$$

- Nucleon's relativistic electromagnetic current:

$$\begin{aligned} J_\mu(P', P) &= ie \bar{u}_p(P') \Lambda_\mu(Q, P) u_p(P), \quad Q = P' - P \\ &= ie \bar{u}_p(P') \left(\gamma_\mu F_1(Q^2) + \frac{1}{2M} \sigma_{\mu\nu} Q_\nu F_2(Q^2) \right) u_p(P) \end{aligned}$$

$$G_E(Q^2) = F_1(Q^2) - \frac{Q^2}{4M^2} F_2(Q^2), \quad G_M(Q^2) = F_1(Q^2) + F_2(Q^2).$$

Point-particle: $F_2 \equiv 0 \Rightarrow G_E \equiv G_M$



Proton Form Factors, Reprise



First

Contents

Back


Conclusion

Proton Form Factors, Reprise

- SLAC and JLab have Measured Ratio of Proton's Electric and Magnetic Form Factors




Proton Form Factors, Reprise




Proton Form Factors, Reprise

- Walker *et al.*, Phys. Rev. D **49**, 5671 (1994). (SLAC)
- Jones *et al.*, JLab Hall A Collaboration, Phys. Rev. Lett. **84**, 1398 (2000)
- Gayou, *et al.*, Phys. Rev. C **64**, 038202 (2001)
- Gayou, *et al.*, JLab Hall A Collaboration, Phys. Rev. Lett. **88** 092301 (2002)




Proton Form Factors, Reprise

- If JLab Correct, then Completely Unexpected Result: In the Proton
 - On Relativistic Domain
 - Distribution of Quark-Charge Not Equal
 - Distribution of Quark-Current!



Modern Miracles in Hadron Physics



First


Contents

Back

Conclusion

Modern Miracles in Hadron Physics

- proton = three constituent quarks



Modern Miracles in Hadron Physics

- proton = three constituent quarks
- $M_{\text{proton}} \approx 1 \text{ GeV}$




Modern Miracles in Hadron Physics

- proton = three constituent quarks
- $M_{\text{proton}} \approx 1 \text{ GeV}$
- guess $M_{\text{constituent-quark}} \approx \frac{1 \text{ GeV}}{3} \approx 350 \text{ MeV}$



Modern Miracles in Hadron Physics

- proton = three constituent **quarks**
- $M_{\text{proton}} \approx 1 \text{ GeV}$
- guess $M_{\text{constituent-quark}} \approx \frac{1 \text{ GeV}}{3} \approx 350 \text{ MeV}$
- pion =
constituent **quark** + constituent **antiquark**



Modern Miracles in Hadron Physics

- proton = three constituent **quarks**
- $M_{\text{proton}} \approx 1 \text{ GeV}$
- guess $M_{\text{constituent-quark}} \approx \frac{1 \text{ GeV}}{3} \approx 350 \text{ MeV}$
- pion =
constituent **quark** + constituent **antiquark**
- guess $M_{\text{pion}} \approx 2 \times \frac{M_{\text{proton}}}{3} \approx 700 \text{ MeV}$



Modern Miracles in Hadron Physics

- proton = three constituent **quarks**
- $M_{\text{proton}} \approx 1 \text{ GeV}$
- guess $M_{\text{constituent-quark}} \approx \frac{1 \text{ GeV}}{3} \approx 350 \text{ MeV}$
- pion =
constituent **quark** + constituent **antiquark**
- guess $M_{\text{pion}} \approx 2 \times \frac{M_{\text{proton}}}{3} \approx 700 \text{ MeV}$
- **WRONG** $M_{\text{pion}} = 140 \text{ MeV}$



Modern Miracles in Hadron Physics

- proton = three constituent **quarks**
- $M_{\text{proton}} \approx 1 \text{ GeV}$
- guess $M_{\text{constituent-quark}} \approx \frac{1 \text{ GeV}}{3} \approx 350 \text{ MeV}$
- pion =
constituent **quark** + constituent **antiquark**
- guess $M_{\text{pion}} \approx 2 \times \frac{M_{\text{proton}}}{3} \approx 700 \text{ MeV}$
- **WRONG** $M_{\text{pion}} = 140 \text{ MeV}$
- Another meson:
..... $M_{\rho} = 770 \text{ MeV}$ **No Surprises Here**



Modern Miracles in Hadron Physics

- proton = three constituent **quarks**
- $M_{\text{proton}} \approx 1 \text{ GeV}$
- guess $M_{\text{constituent-quark}} \approx \frac{1 \text{ GeV}}{3} \approx 350 \text{ MeV}$
- pion =
constituent **quark** + constituent **antiquark**
- guess $M_{\text{pion}} \approx 2 \times \frac{M_{\text{proton}}}{3} \approx 700 \text{ MeV}$
- **WRONG** $M_{\text{pion}} = 140 \text{ MeV}$
- What is “**wrong**” with the pion?



Dichotomy of the Pion





Dichotomy of the Pion

- How does one make an **almost massless** particle
..... from two **massive** constituent-quarks?





Dichotomy of the Pion

- How does one make an **almost massless** particle from two **massive** constituent-quarks?
- **Not Allowed** to do it by **fine-tuning** a potential

Must exhibit $m_\pi^2 \propto m_q$

Current Algebra ... 1968





Dichotomy of the Pion

- How does one make an **almost massless** particle from two **massive** constituent-quarks?
- **Not Allowed** to do it by **fine-tuning** a potential
Must exhibit $m_\pi^2 \propto m_q$

Current Algebra ... 1968

The **correct understanding** of pion observables;
e.g. **mass**, **decay constant** and **form factors**,
requires an approach to contain a **well-defined** and
valid chiral limit, and an **accurate realisation** of
dynamical chiral symmetry breaking.





Dichotomy of the Pion

- How does one make an **almost massless** particle from two **massive** constituent-quarks?
- **Not Allowed** to do it by **fine-tuning** a potential

Must exhibit $m_\pi^2 \propto m_q$

Current Algebra ... 1968

The **correct understanding** of pion observables;
e.g. **mass**, **decay constant** and **form factors**,
requires an approach to contain a **well-defined** and
valid chiral limit, and an **accurate realisation** of
dynamical chiral symmetry breaking.

- **Requires** detailed understanding of Connection between **Current-quark** and **Constituent-quark** masses





Dichotomy of the Pion

- How does one make an **almost massless** particle from two **massive** constituent-quarks?
- Not Allowed to do it by **fine-tuning** a potential
Must exhibit $m_\pi^2 \propto m_q$

Current Algebra ... 1968

The **correct understanding** of pion observables;
e.g. **mass**, **decay constant** and **form factors**,
requires an approach to contain a **well-defined** and
valid chiral limit, and an **accurate realisation** of
dynamical chiral symmetry breaking.


- **Requires** detailed understanding of Connection
between **Current-quark** and **Constituent-quark**
masses

Using DSEs,
we've provided this.
Dyson-Schwinger Equations – Achievements and Challenges
Peking University, Beijing, China, 13-17 August, 2007



QCD's Emergent Phenomena






QCD's Emergent Phenomena

- Complex behaviour arises from apparently simple rules






QCD's Emergent Phenomena

- Complex behaviour arises from apparently simple rules
- Quark and Gluon Confinement
 - No matter how hard one strikes the proton, one cannot liberate an individual quark or gluon





QCD's Emergent Phenomena

- Complex behaviour arises from apparently simple rules
- Quark and Gluon Confinement
 - No matter how hard one strikes the proton, one cannot liberate an individual quark or gluon
- Dynamical Chiral Symmetry Breaking
 - Very unnatural pattern of bound state masses






QCD's Emergent Phenomena

- Complex behaviour arises from apparently simple rules
- Quark and Gluon Confinement
 - No matter how hard one strikes the proton, one cannot liberate an individual quark or gluon
- Dynamical Chiral Symmetry Breaking
 - Very unnatural pattern of bound state masses
- Neither of these phenomena is apparent in QCD's Lagrangian yet they are the dominant determining characteristics of real-world QCD.





QCD's Emergent Phenomena

- Complex behaviour arises from apparently simple rules
- Quark and Gluon Confinement
 - No matter how hard one strikes the proton, one cannot liberate an individual quark or gluon
- Dynamical Chiral Symmetry Breaking
 - Very unnatural pattern of bound state masses
- Neither of these phenomena is apparent in QCD's Lagrangian yet they are the dominant determining characteristics of real-world QCD.
- NSAC – Understanding these phenomena is one of the greatest intellectual challenges in physics



What's the Problem?



First

Contents

Back

Conclusion

What's the Problem?

- Must calculate the hadron's *wave function*
 - Can't be done using perturbation theory



What's the Problem?

- Must calculate the hadron's *wave function*
 - Can't be done using perturbation theory
 - So what? Same is true of hydrogen atom



What's the Problem?

- Must calculate the hadron's *wave function*
 - Can't be done using perturbation theory
 - So what? Same is true of hydrogen atom
- Differences



What's the Problem?

- Must calculate the hadron's *wave function*
 - Can't be done using perturbation theory
 - So what? Same is true of hydrogen atom
- Differences
 - Here relativistic effects are crucial
 - *virtual particles*

Quintessence of Relativistic Quantum Field Theory



What's the Problem?

- Must calculate the hadron's *wave function*
 - Can't be done using perturbation theory
 - So what? Same is true of hydrogen atom
- Differences
 - Here relativistic effects are crucial
 - *virtual particles*
- Interaction between quarks – the *Interquark Potential* –
 - Unknown
 - throughout $> 98\%$ of the pion's/proton's volume




What's the Problem?

- Must calculate the hadron's *wave function*
 - Can't be done using perturbation theory
 - So what? Same is true of hydrogen atom
- Differences
 - Here relativistic effects are crucial
 - *virtual particles*
Quintessence of Relativistic Quantum Field Theory
 - Interaction between quarks – the *Interquark Potential* – *Unknown*
throughout $> 98\%$ of the pion's/proton's volume
 - Determination of hadrons's wave function requires
ab initio nonperturbative solution
of fully-fledged relativistic quantum field theory



What's the Problem?


- Must calculate the hadron's *wave function*
 - Can't be done using perturbation theory
 - So what? Same is true of hydrogen atom
- Determination of hadron's wave function requires *ab initio* nonperturbative solution of fully-fledged relativistic quantum field theory
- Modern Physics & Mathematics
 - Still quite some way from being able to do that




Intranucleon Interaction



Intranucleon Interaction



Intranucleon Interaction



98% of the volume



Dyson-Schwinger Equations



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
..... Materially Reduces Model Dependence



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
 - Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
..... Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
 - Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons
 - Qualitative and Quantitative Importance of:
 - Dynamical Chiral Symmetry Breaking
 - Generation of fermion mass from *nothing*
 - Quark & Gluon Confinement
 - Coloured objects not detected, not detectable?



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
 - Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons
 - Qualitative and Quantitative Importance of:
 - Dynamical Chiral Symmetry Breaking
 - Generation of fermion mass from *nothing*
 - Quark & Gluon Confinement
 - Coloured objects not detected, not detectable?
 - ⇒ Understanding InfraRed (long-range)
 - behaviour of $\alpha_s(Q^2)$



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
 - Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons
 - Qualitative and Quantitative Importance of:
 - Dynamical Chiral Symmetry Breaking
 - Generation of fermion mass from *nothing*
 - Quark & Gluon Confinement
 - Coloured objects not detected, not detectable?
 - Method yields Schwinger Functions \equiv Propagators



Dyson-Schwinger Equations


- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
 - Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons
 - Qualitative and Quantitative Importance of:
 - Dynamical Chiral Symmetry Breaking
 - Generation of fermion mass from *nothing*
 - Quark & Gluon Confinement
 - Coloured objects not detected, not detectable?

Cross-Sections built from Schwinger Functions




Persistent Challenge





Persistent Challenge

- Infinitely Many Coupled Equations





Persistent Challenge

- Infinitely Many Coupled Equations
 - Solutions are **Schwinger Functions**
(Euclidean **Green Functions**)





Persistent Challenge

- Infinitely Many Coupled Equations
 - Solutions are Schwinger Functions
(Euclidean **Green** Functions)
 - Not all are Schwinger functions are experimentally observable but **all** are same VEVs measured in Lattice-QCD simulations . . . opportunity for comparisons at pre-experimental level . . . cross-fertilisation





Persistent Challenge

- Infinitely Many Coupled Equations
 - Solutions are Schwinger Functions
(Euclidean **Green** Functions)
- Coupling between equations **necessitates** truncation
 - Weak coupling expansion \Rightarrow Perturbation Theory





Persistent Challenge

- Infinitely Many Coupled Equations
 - Solutions are Schwinger Functions
(Euclidean **Green** Functions)
- Coupling between equations **necessitates** truncation
 - Weak coupling expansion \Rightarrow Perturbation Theory
Not useful for the nonperturbative problems
in which we're interested





Persistent Challenge

- Infinitely Many Coupled Equations
 - Solutions are Schwinger Functions
(Euclidean **Green** Functions)
- There is at least one systematic nonperturbative, symmetry-preserving truncation scheme
H.J. Munczek Phys. Rev. D **52** (1995) 4736
Dynamical chiral symmetry breaking, Goldstone's theorem and the consistency of the Schwinger-Dyson and Bethe-Salpeter Equations
A. Bender, C. D. Roberts and L. von Smekal, Phys. Lett. B **380** (1996) 7
Goldstone Theorem and Diquark Confinement Beyond Rainbow Ladder Approximation





Persistent Challenge

- Infinitely Many Coupled Equations
 - Solutions are **Schwinger Functions** (Euclidean **Green Functions**)
- There is at least one **systematic nonperturbative, symmetry-preserving truncation scheme**
- Has Enabled Proof of **EXACT** Results in QCD





Persistent Challenge

- Infinitely Many Coupled Equations
 - Solutions are Schwinger Functions
(Euclidean **Green** Functions)
- There is at least one **systematic nonperturbative, symmetry-preserving** truncation scheme
- Has Enabled Proof of **EXACT** Results in QCD
- And Formulation of Practical Phenomenological Tool to
 - Illustrate Exact Results





Persistent Challenge

- Infinitely Many Coupled Equations
 - Solutions are Schwinger Functions
(Euclidean **Green** Functions)
- There is at least one **systematic nonperturbative, symmetry-preserving** truncation scheme
- Has Enabled Proof of **EXACT** Results in QCD
- And Formulation of Practical Phenomenological Tool to
 - Make Predictions with Readily Quantifiable Errors




Dressed-Quark Propagator



Dressed-Quark Propagator

$$S(p) = \frac{Z(p^2)}{i\gamma \cdot p + M(p^2)}$$




Gap Equation




Dressed-Quark Propagator

$$S(p) = \frac{Z(p^2)}{i\gamma \cdot p + M(p^2)}$$




Gap Equation

- Gap Equation's Kernel Enhanced on IR domain
⇒ IR Enhancement of $M(p^2)$



Dressed-Quark Propagator

$$S(p) = \frac{Z(p^2)}{i\gamma \cdot p + M(p^2)}$$




Gap Equation

- Gap Equation's Kernel Enhanced on **IR domain**
⇒ **IR Enhancement of $M(p^2)$**

- Euclidean Constituent–Quark Mass: M_f^E : $p^2 = M(p^2)^2$

flavour	u/d	s	c	b
$\frac{M_f^E}{m_\zeta}$	$\sim 10^2$	~ 10	~ 1.5	~ 1.1



Dressed-Quark Propagator



- Longstanding Prediction of Dyson-Schwinger Equation Studies



Dressed-Quark Propagator




- Longstanding Prediction of Dyson-Schwinger Equation Studies
 - E.g., *Dyson-Schwinger equations and their application to hadronic physics*,
C. D. Roberts and
A. G. Williams,
Prog. Part. Nucl. Phys.
33 (1994) 477



Dressed-Quark Propagator

DO YOU
THINK KEN'S
CONSTIPATION
WILL END
HAPPILY?

The ending is
unimportant; what
matters most is
the sheer drama
of his difficult
and lonely
situation.



[167]

- Long used as basis for efficacious hadron physics phenomenology




- Longstanding Prediction of Dyson-Schwinger Equation Studies
 - E.g., *Dyson-Schwinger equations and their application to hadronic physics*,
C. D. Roberts and
A. G. Williams,
Prog. Part. Nucl. Phys.
33 (1994) 477

Dressed-Quark Propagator

DO YOU
THINK KEN'S
CONSTIPATION
WILL END
HAPPILY?



The ending is
unimportant; what
matters most is
the sheer drama
of his difficult
and lonely
situation.



- Long used as basis for efficacious hadron physics phenomenology
 - *Electromagnetic pion form-factor and neutral pion decay width,*
C. D. Roberts,
Nucl. Phys. A **605**
(1996) 475





Dressed-Quark Propagator

 $M(p)$  $Z(p)$ 

2002

Dressed-Quark Propagator

 $M(p)$  $Z(p)$ 



“data:” Quenched Lattice Meas.

– Bowman, Heller, Leinweber, Williams: [he-lat/0209129](https://arxiv.org/abs/hep-lat/0209129)



2002

Dressed-Quark Propagator

 $M(p)$  $Z(p)$ 



“data:” Quenched Lattice Meas.

- Bowman, Heller, Leinweber, Williams: [he-lat/0209129](https://arxiv.org/abs/hep-lat/0209129)
- current-quark masses: 30 MeV, 50 MeV, 100 MeV



2002

Dressed-Quark Propagator

 $M(p)$  $Z(p)$ 



- “*data*:” Quenched Lattice Meas.
 - Bowman, Heller, Leinweber, Williams: [he-lat/0209129](#)
current-quark masses: 30 MeV, 50 MeV, 100 MeV
- *Curves*: Quenched DSE Cal.
 - Bhagwat, Pichowsky, Roberts, Tandy [nu-th/0304003](#)



Dressed-Quark Propagator



2002

 $M(p)$  $Z(p)$ 

“data:” Quenched Lattice Meas.

- Bowman, Heller, Leinweber, Williams: [he-lat/0209129](#)
current-quark masses: 30 MeV, 50 MeV, 100 MeV

Curves: Quenched DSE Cal.

- Bhagwat, Pichowsky, Roberts, Tandy [nu-th/0304003](#)

Linear extrapolation of lattice data to chiral limit is inaccurate



QCD & Interaction Between Light-Quarks

- Kernel of Gap Equation: $D_{\mu\nu}(p - q) \Gamma_\nu(q)$
Dressed-gluon propagator and dressed-quark-gluon vertex
- Reliable DSE studies of Dressed-gluon propagator:
 - R. Alkofer and L. von Smekal, *The infrared behavior of QCD Green's functions . . .*, Phys. Rept. **353**, 281 (2001).



QCD & Interaction Between Light-Quarks

- Kernel of Gap Equation: $D_{\mu\nu}(p - q) \Gamma_\nu(q)$
Dressed-gluon propagator and dressed-quark-gluon vertex
- Reliable DSE studies of Dressed-gluon propagator:
 - R. Alkofer and L. von Smekal, *The infrared behavior of QCD Green's functions . . .*, Phys. Rept. **353**, 281 (2001).
- Dressed-gluon propagator – lattice-QCD simulations confirm that behaviour:
 - D. B. Leinweber, J. I. Skullerud, A. G. Williams and C. Parrinello [UKQCD Collaboration], *Asymptotic scaling and infrared behavior of the gluon propagator*, Phys. Rev. D **60**, 094507 (1999) [Erratum-ibid. D **61**, 079901 (2000)].
- Exploratory DSE and lattice-QCD studies of dressed-quark-gluon vertex




Dressed-gluon Propagator

- $D_{\mu\nu}(k) = \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) \frac{Z(k^2)}{k^2}$

- Suppression means \exists IR gluon mass-scale $\approx 1 \text{ GeV}$

- Naturally, this scale has the same origin as Λ_{QCD}




Dressed-gluon Propagator

- $D_{\mu\nu}(k) = \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) \frac{Z(k^2)}{k^2}$

- Suppression means \exists IR gluon mass-scale $\approx 1 \text{ GeV}$

- Naturally, this scale has the same origin as Λ_{QCD}




Dressed-gluon Propagator

- $D_{\mu\nu}(k) = \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) \frac{Z(k^2)}{k^2}$

- Suppression means \exists IR gluon mass-scale $\approx 1 \text{ GeV}$

- Naturally, this scale has the same origin as Λ_{QCD}



Critical Mass for Chiral Expansion

Dynamical chiral symmetry breaking and a critical mass

Lei Chang, Yu-Xin Liu, Mandar S. Bhagwat, Craig D. Roberts
and Stewart V. Wright . . . nucl-th/0605058
Phys. Rev. C 75 (2007) 015201 (8 pages)




Critical Mass for Chiral Expansion

- Realistic models of QCD's gap equation simultaneously admit two inequivalent DCSB solutions & solution connected with realisation of chiral symmetry in Wigner mode.



Critical Mass for Chiral Expansion


- Realistic models of QCD's gap equation simultaneously admit two inequivalent DCSB solutions & solution connected with realisation of chiral symmetry in Wigner mode.
- Wigner solution and one DCSB solution are destabilised by current-quark mass & both disappear when that mass exceeds a critical value, m_{cr}



The zeros of $G(M)$ characterise the different solutions of the gap equation.
Solid curve: obtained with $m^{\text{bm}} = 0$, in which case $G(M)$ is odd under $M \rightarrow -M$; long-dashed curve: $m^{\text{bm}} = 0.01$; short-dashed curve: $m^{\text{bm}} = m_{\text{cr}}^{\text{bm}} = 0.033$; dotted curve: $m^{\text{bm}} = 0.05$.

Critical Mass for Chiral Expansion

- Realistic models of QCD's gap equation simultaneously admit two inequivalent DCSB solutions & solution connected with realisation of chiral symmetry in Wigner mode.
- m_{cr} also bounds domain on which surviving DCSB solution possesses a chiral expansion: $m_{\text{cr}} = \lim_{n \rightarrow \infty} \left(\frac{1}{|a_n|} \right)^{1/n}$



Critical Mass for Chiral Expansion

- Realistic models of QCD's gap equation simultaneously admit two inequivalent DCSB solutions & solution connected with realisation of chiral symmetry in Wigner mode.
- m_{cr} also bounds domain on which surviving DCSB solution possesses a chiral expansion: $m_{\text{cr}} = \lim_{n \rightarrow \infty} \left(\frac{1}{|a_n|} \right)^{1/n}$
- m_{cr} is therefore an upper bound on the domain within which a perturbative expansion in the current-quark mass around the chiral limit is uniformly valid for physical quantities.



Critical Mass for Chiral Expansion

- Realistic models of QCD's gap equation simultaneously admit two inequivalent DCSB solutions & solution connected with realisation of chiral symmetry in Wigner mode.
- m_{cr} also bounds domain on which surviving DCSB solution possesses a chiral expansion: $m_{\text{cr}} = \lim_{n \rightarrow \infty} \left(\frac{1}{|a_n|} \right)^{1/n}$
- m_{cr} is therefore an upper bound on the domain within which a perturbative expansion in the current-quark mass around the chiral limit is uniformly valid for physical quantities.
- For a pseudoscalar meson constituted of equal mass current-quarks, it corresponds to a mass $\textcolor{red}{m_{0^-}^{\text{cr}}} \sim 0.45 \text{ GeV}$.



Critical Mass for Chiral Expansion

- Realistic models of QCD's gap equation simultaneously admit two inequivalent DCSB solutions & solution connected with realisation of chiral symmetry in Wigner mode.
- m_{cr} also bounds domain on which surviving DCSB solution possesses a chiral expansion: $m_{\text{cr}} = \lim_{n \rightarrow \infty} \left(\frac{1}{|a_n|} \right)^{1/n}$
- m_{cr} is therefore an upper bound on the domain within which a perturbative expansion in the current-quark mass around the chiral limit is uniformly valid for physical quantities.
- For a pseudoscalar meson constituted of equal mass current-quarks, it corresponds to a mass $m_{0^-}^{\text{cr}} \sim 0.45 \text{ GeV}$.
- Entails lattice-QCD simulations *must* have many results at $m_\pi < m_{0^-}^{\text{cr}} \sim 0.45 \text{ GeV}$ for reliable extrapolation via EFT.




Critical Mass for Chiral Expansion

- Realistic models of QCD's gap equation simultaneously admit two inequivalent DCSB solutions & solution connected with realisation of chiral symmetry in Wigner mode.
- The two DCSB solutions of the gap equation enable a valid definition of $\langle \bar{q}q \rangle$ in the presence of a nonzero current-mass.



Critical Mass for Chiral Expansion

- Realistic models of QCD's gap equation simultaneously admit two inequivalent DCSB solutions & solution connected with realisation of chiral symmetry in Wigner mode.
- The two DCSB solutions of the gap equation enable a valid definition of $\langle \bar{q}q \rangle$ in the presence of a nonzero current-mass.
- The behaviour of this condensate indicates that the essentially dynamical component of chiral symmetry breaking decreases with increasing current-quark mass.



Constituent-quark σ -term

- Impact of Dynamical chiral symmetry breaking . . . exhibited via constituent-quark σ -term

$$\sigma_f := m_f(\zeta) \frac{\partial M_f^E}{\partial m_f(\zeta)}, \quad (M^E)^2 := s \mid s = M(s)^2.$$



Constituent-quark σ -term

- Impact of Dynamical chiral symmetry breaking . . . exhibited via constituent-quark σ -term

$$\sigma_f := m_f(\zeta) \frac{\partial M_f^E}{\partial m_f(\zeta)}, \quad (M^E)^2 := s \mid s = M(s)^2.$$

- Renormalisation-group-invariant and determined from solutions of the gap equation



Constituent-quark σ -term

- Impact of Dynamical chiral symmetry breaking . . . exhibited via constituent-quark σ -term

$$\sigma_f := m_f(\zeta) \frac{\partial M_f^E}{\partial m_f(\zeta)}, \quad (M^E)^2 := s \mid s = M(s)^2.$$

- Unambiguous probe of impact of explicit chiral symmetry breaking on the mass function



Constituent-quark σ -term

- Impact of Dynamical chiral symmetry breaking . . . exhibited via constituent-quark σ -term

$$\sigma_f := m_f(\zeta) \frac{\partial M_f^E}{\partial m_f(\zeta)}, \quad (M^E)^2 := s \mid s = M(s)^2.$$


- Ratio
$$\frac{\sigma_f}{M_f^E} = \frac{\text{EXPLICIT}}{\text{EXPLICIT} + \text{DYNAMICAL}}$$
measures effect of **EXPLICIT** chiral symmetry breaking on dressed-quark mass-function
cf. **SUM** of effects of **EXPLICIT AND DYNAMICAL CHIRAL SYMMETRY BREAKING**



Constituent-quark σ -term

- Impact of Dynamical chiral symmetry breaking . . . exhibited via constituent-quark σ -term

$$\sigma_f := m_f(\zeta) \frac{\partial M_f^E}{\partial m_f(\zeta)}, \quad (M^E)^2 := s \mid s = M(s)^2.$$




Argonne
NATIONAL
LABORATORY

Constituent-quark σ -term

- Impact of Dynamical chiral symmetry breaking . . . exhibited via constituent-quark σ -term

$$\sigma_f := m_f(\zeta) \frac{\partial M_f^E}{\partial m_f(\zeta)}, \quad (M^E)^2 := s \mid s = M(s)^2.$$

Obvious: ratio vanishes for light-quarks because magnitude of their constituent-mass owes primarily to DCSB. On the other hand, for heavy-quarks it approaches one.




Constituent-quark σ -term

- Impact of Dynamical chiral symmetry breaking . . . exhibited via constituent-quark σ -term

$$\sigma_f := m_f(\zeta) \frac{\partial M_f^E}{\partial m_f(\zeta)}, \quad (M^E)^2 := s \mid s = M(s)^2.$$

Essentially dynamical component of chiral symmetry breaking, and manifestation in all its order parameters, vanishes with increasing current-quark mass





Hadrons

- Established understanding of two- and three-point functions





Hadrons

- Established understanding of two- and three-point functions
- What about bound states?





Hadrons


- Without bound states,
Comparison with experiment is
impossible



- Without bound states,
Comparison with experiment is
impossible
- They appear as pole contributions
to $n \geq 3$ -point colour-singlet
Schwinger functions




- Without bound states,
Comparison with experiment is
impossible
- Bethe-Salpeter Equation



QFT Generalisation of Lippmann-Schwinger Equation.



- Without bound states,
Comparison with experiment is
impossible
- Bethe-Salpeter Equation




QFT Generalisation of Lippmann-Schwinger Equation.

- What is the kernel, K ?



- Without bound states,
Comparison with experiment is
impossible
- Bethe-Salpeter Equation



QFT Generalisation of Lippmann-Schwinger Equation.

- What is the kernel, K ?

or

What is the Long-Range Potential?



First


Contents

Back

Conclusion

What is the Long-Range Potential?

Bush Urges Nation To Be Quiet For A Minute While He Tries To Think



In a televised address to the nation, Bush called for "a little peace and quiet."



Bethe-Salpeter Kernel



Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i\gamma_5 + \frac{1}{2} \lambda_f^l i\gamma_5 \mathcal{S}^{-1}(k_-)$$

$$-M_\zeta i\Gamma_5^l(k; P) - i\Gamma_5^l(k; P) M_\zeta$$

QFT Statement of Chiral Symmetry



Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P)$$

$$= \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5 \mathcal{S}^{-1}(k_-)$$

$$- M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE



Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P)$$

$$= \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5 \mathcal{S}^{-1}(k_-)$$

$$- M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE

Kernels must be **intimately** related



Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P)$$

$$= \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5$$

$$\mathcal{S}^{-1}(k_-)$$

$$- M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE

Kernels must be **intimately** related

- Relation **must** be preserved by truncation



Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i\gamma_5 + \frac{1}{2} \lambda_f^l i\gamma_5 \mathcal{S}^{-1}(k_-)$$
$$- M_\zeta i\Gamma_5^l(k; P) - i\Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE

Kernels must be **intimately** related

- Relation **must** be preserved by truncation
- **Nontrivial** constraint





Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i\gamma_5 + \frac{1}{2} \lambda_f^l i\gamma_5 \mathcal{S}^{-1}(k_-)$$
$$- M_\zeta i\Gamma_5^l(k; P) - i\Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE


Satisfies DSE

Kernels must be **intimately** related

- Relation **must** be preserved by truncation
- **Failure** \Rightarrow Explicit Violation of QCD's Chiral Symmetry




Pseudoscalar Mesons?




Pseudoscalar Mesons?

Can a bound-state of massive constituents truly be massless ... **without** fine-tuning?



Radial Excitations & Chiral Symmetry



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \quad m_H^2 = - \rho_{\zeta}^H \mathcal{M}_H$$

- Mass² of pseudoscalar hadron



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \quad m_H^2 = - \quad \rho_{\zeta}^H \quad \mathcal{M}_H$$

$$\mathcal{M}_H := \text{tr}_{\text{flavour}} \left[\mathcal{M}_{(\mu)} \left\{ T^H, (T^H)^t \right\} \right] = m_{q_1} + m_{q_2}$$

- Sum of constituents' current-quark masses
- e.g., $T^{K^+} = \frac{1}{2} (\lambda^4 + i\lambda^5)$




Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H p_\mu = Z_2 \int_q^\Lambda \frac{1}{2} \text{tr} \left\{ (T^H)^t \gamma_5 \gamma_\mu \boxed{\mathcal{S}(q_+) \Gamma_H(q; P) \mathcal{S}(q_-)} \right\}$$

$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$

- Pseudovector projection of BS wave function at $x = 0$
- Pseudoscalar meson's leptonic decay constant




Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \quad m_H^2 = - \rho_{\zeta}^H \mathcal{M}_H$$

$$i\rho_{\zeta}^H = Z_4 \int_q^{\Lambda} \frac{1}{2} \text{tr} \left\{ (T^H)^t \gamma_5 \mathcal{S}(q_+) \Gamma_H(q; P) \mathcal{S}(q_-) \right\}$$

- Pseudoscalar projection of BS wave function at $x = 0$



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Light-quarks; i.e., $m_q \sim 0$

- $f_H \rightarrow f_H^0$ & $\rho_\zeta^H \rightarrow \frac{-\langle \bar{q}q \rangle_\zeta^0}{f_H^0}$, Independent of m_q

Hence $m_H^2 = \frac{-\langle \bar{q}q \rangle_\zeta^0}{(f_H^0)^2} m_q$... GMOR relation, a corollary



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Light-quarks; i.e., $m_q \sim 0$

- $f_H \rightarrow f_H^0$ & $\rho_\zeta^H \rightarrow \frac{-\langle \bar{q}q \rangle_\zeta^0}{f_H^0}$, Independent of m_q

Hence
$$m_H^2 = \frac{-\langle \bar{q}q \rangle_\zeta^0}{(f_H^0)^2} m_q \dots \text{GMOR relation, a corollary}$$

- Heavy-quark + light-quark

$$\Rightarrow f_H \propto \frac{1}{\sqrt{m_H}} \text{ and } \rho_\zeta^H \propto \sqrt{m_H}$$

Hence,
$$m_H \propto m_q$$

... QCD Proof of Potential Model result

Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \quad m_H^2 = - \quad \rho_\zeta^H \quad \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson,
 $m_{\pi_n \neq 0}^2 > m_{\pi_n = 0}^2 = 0$, in chiral limit



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \quad m_H^2 = - \quad \rho_\zeta^H \quad \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson,
 $m_{\pi_n \neq 0}^2 > m_{\pi_n = 0}^2 = 0$, in chiral limit
- $\Rightarrow f_H = 0$
ALL pseudoscalar mesons except $\pi(140)$ in chiral limit



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson,
 $m_{\pi_n \neq 0}^2 > m_{\pi_n = 0}^2 = 0$, in chiral limit
- $\Rightarrow f_H = 0$
ALL pseudoscalar mesons except $\pi(140)$ in chiral limit
- Dynamical Chiral Symmetry Breaking
 - Goldstone’s Theorem –impacts upon *every* pseudoscalar meson



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- ρ_H \Rightarrow finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- ρ_H \Rightarrow finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson,
 $m_{\pi_n \neq 0}^2 > m_{\pi_n = 0}^2 = 0$, in chiral limit



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \quad m_H^2 = - \quad \rho_\zeta^H \quad \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson,
 $m_{\pi_n \neq 0}^2 > m_{\pi_n = 0}^2 = 0$, in chiral limit
- $\Rightarrow f_H = 0$
ALL pseudoscalar mesons except $\pi(140)$ in chiral limit



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H \ m_H^2 = - \rho_\zeta^H \ \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson,
 $m_{\pi_n \neq 0}^2 > m_{\pi_n = 0}^2 = 0$, in chiral limit
- $\Rightarrow f_H = 0$
ALL pseudoscalar mesons except $\pi(140)$ in chiral limit
- Dynamical Chiral Symmetry Breaking
 - Goldstone’s Theorem –impacts upon *every* pseudoscalar meson



Radial Excitations

& Lattice-QCD

McNeile and Michael
he-la/0607032



First

Contents

Back

Conclusion

Radial Excitations & Lattice-QCD

McNeile and Michael
he-la/0607032

- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.



Radial Excitations & Lattice-QCD

McNeile and Michael
he-la/0607032



- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.
- CLEO: $\tau \rightarrow \pi(1300) + \nu_\tau$
 $\Rightarrow f_{\pi_1} < 8.4 \text{ MeV}$
Diehl & Hiller
he-ph/0105194



Radial Excitations & Lattice-QCD

McNeile and Michael
he-la/0607032

- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.
- CLEO: $\tau \rightarrow \pi(1300) + \nu_\tau$
 $\Rightarrow f_{\pi_1} < 8.4 \text{ MeV}$
Diehl & Hiller
he-ph/0105194
- Lattice-QCD check:
 $16^3 \times 32$,
 $a \sim 0.1 \text{ fm}$,
two-flavour, unquenched
 $\Rightarrow \frac{f_{\pi_1}}{f_\pi} = 0.078 (93)$



Radial Excitations & Lattice-QCD

McNeile and Michael
he-la/0607032


- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.

- CLEO: $\tau \rightarrow \pi(1300) + \nu_\tau$
 $\Rightarrow f_{\pi_1} < 8.4 \text{ MeV}$

Diehl & Hiller
he-ph/0105194

- Lattice-QCD check:
 $16^3 \times 32$,
 $a \sim 0.1 \text{ fm}$,
two-flavour, unquenched

$$\Rightarrow \frac{f_{\pi_1}}{f_\pi} = 0.078 (93)$$



- Full ALPHA formulation is required to see suppression, because PCAC relation is at the heart of the conditions imposed for improvement (determining coefficients of irrelevant operators)


- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.

- CLEO: $\tau \rightarrow \pi(1300) + \nu_\tau$
 $\Rightarrow f_{\pi_1} < 8.4 \text{ MeV}$

Diehl & Hiller
he-ph/0105194

- Lattice-QCD check:
 $16^3 \times 32$,
 $a \sim 0.1 \text{ fm}$,
two-flavour, unquenched

$$\Rightarrow \frac{f_{\pi_1}}{f_\pi} = 0.078 (93)$$



- The suppression of f_{π_1} is a useful benchmark that can be used to tune and validate lattice QCD techniques that try to determine the properties of excited states mesons.



Radial Excitations



Radial Excitations

- Spectrum contains 3 pseudoscalars [$I^G(J^P)L = 1^-(0^-)S$]

masses below 2 GeV: $\pi(140)$; $\pi(1300)$; and $\pi(1800)$



Radial Excitations

- Spectrum contains 3 pseudoscalars [$I^G(J^P)L = 1^-(0^-)S$]
masses below 2 GeV: $\pi(140)$; $\pi(1300)$; and $\pi(1800)$
- The Pion
- Constituent-Q Model: 1st three members of $n\ ^1S_0$ trajectory; i.e., ground state plus radial excitations?



Radial Excitations

- Spectrum contains 3 pseudoscalars [$I^G(J^P)L = 1^-(0^-)S$]
masses below 2 GeV: $\pi(140)$; $\pi(1300)$; and $\pi(1800)$
- The Pion
- Constituent-Q Model: 1st three members of $n\ ^1S_0$ trajectory; i.e., ground state plus radial excitations?
- But $\pi(1800)$ is narrow ($\Gamma = 207 \pm 13$) & decay pattern might indicate some “flux tube angular momentum” content:
 $S_{\bar{Q}Q} = 1 \oplus L_F = 1 \Rightarrow J = 0$
& $L_F = 1 \Rightarrow ^3S_1 \oplus ^3S_1 (\bar{Q}Q)$ decays suppressed?



Radial Excitations

- Spectrum contains 3 pseudoscalars [$I^G(J^P)L = 1^-(0^-)S$]
masses below 2 GeV: $\pi(140)$; $\pi(1300)$; and $\pi(1800)$
- The Pion
- Constituent-Q Model: 1st three members of $n\ ^1S_0$ trajectory; i.e., ground state plus radial excitations?
- But $\pi(1800)$ is narrow ($\Gamma = 207 \pm 13$) & decay pattern might indicate some “flux tube angular momentum” content:
- Radial excitations & Hybrids & Exotics \Rightarrow Long-range radial wave functions \Rightarrow sensitive to confinement



Radial Excitations

- Spectrum contains 3 pseudoscalars [$I^G(J^P)L = 1^-(0^-)S$]
masses below 2 GeV: $\pi(140)$; $\pi(1300)$; and $\pi(1800)$
- The Pion
- Constituent-Q Model: 1st three members of $n\ ^1S_0$ trajectory; i.e., ground state plus radial excitations?
- But $\pi(1800)$ is narrow ($\Gamma = 207 \pm 13$) & decay pattern might indicate some “flux tube angular momentum” content:
- Radial excitations & Hybrids & Exotics \Rightarrow Long-range radial wave functions \Rightarrow sensitive to confinement
- NSAC Long-Range Plan, 2002: . . . an understanding of confinement “remains one of the greatest intellectual challenges in physics”



but ...

- Orbital angular momentum is not a Poincaré invariant.
However, if absent in a particular frame, it will appear in another frame related via a Poincaré transformation.



but ...

- Nonzero quark orbital angular momentum is thus a necessary outcome of a Poincaré covariant description.



but ...

- Pseudoscalar meson Bethe-Salpeter amplitude

$$\begin{aligned}\chi_{\pi}(k; P) &= \gamma_5 [i\mathcal{E}_{\pi_n}(k; P) + \gamma \cdot P \mathcal{F}_{\pi_n}(k; P) \\ &\quad \gamma \cdot k \, k \cdot P \mathcal{G}_{\pi_n}(k; P) + \sigma_{\mu\nu} k_{\mu} P_{\nu} \mathcal{H}_{\pi_n}(k; P)]\end{aligned}$$



but ...

- Pseudoscalar meson Bethe-Salpeter amplitude

$$\begin{aligned}\chi_{\pi}(k; P) &= \gamma_5 [i\mathcal{E}_{\pi_n}(k; P) + \gamma \cdot P \mathcal{F}_{\pi_n}(k; P) \\ &\quad \gamma \cdot k \, k \cdot P \mathcal{G}_{\pi_n}(k; P) + \sigma_{\mu\nu} k_{\mu} P_{\nu} \mathcal{H}_{\pi_n}(k; P)]\end{aligned}$$

- $J = 0 \dots$ *but* while \mathcal{E} and \mathcal{F} are purely $L = 0$ in the rest frame, the \mathcal{G} and \mathcal{H} terms are associated with $L = 1$. Thus a pseudoscalar meson Bethe-Salpeter wave function *always* contains both S - and P -wave components.



but ...

- $J = 0 \dots$ but while \mathcal{E} and \mathcal{F} are purely $L = 0$ in the rest frame, the \mathcal{G} and \mathcal{H} terms are associated with $L = 1$. Thus a pseudoscalar meson Bethe-Salpeter wave function *always* contains both S - and P -wave components.

Introduce mixing

angle θ_π such that

$$\chi_\pi \sim \cos \theta_\pi |L = 0\rangle$$

$$+ \sin \theta_\pi |L = 1\rangle$$




but ...

- $J = 0 \dots$ but while \mathcal{E} and \mathcal{F} are purely $L = 0$ in the rest frame, the \mathcal{G} and \mathcal{H} terms are associated with $L = 1$. Thus a pseudoscalar meson Bethe-Salpeter wave function *always* contains both S - and P -wave components.

Introduce mixing angle θ_π such that

$$\chi_\pi \sim \cos \theta_\pi |L = 0\rangle + \sin \theta_\pi |L = 1\rangle$$





but ...

- $J = 0 \dots$ but while \mathcal{E} and \mathcal{F} are purely $L = 0$ in the rest frame, the \mathcal{G} and \mathcal{H} terms are associated with $L = 1$. Thus a pseudoscalar meson Bethe-Salpeter wave function *always* contains both S - and P -wave components.

Introduce mixing angle θ_π such that

$$\chi_\pi \sim \cos \theta_\pi |L = 0\rangle + \sin \theta_\pi |L = 1\rangle$$

L is significant in the neighbourhood of the chiral limit, and decreases with increasing current-quark mass.



Charge Neutral Pseudoscalar Mesons

Flavour symmetry breaking and meson masses

Mandar S. Bhagwat, Lei Chang, Yu-Xin Liu, Craig D. Roberts
and Peter C. Tandy ... nucl-th/arXiv:0708.1118



Charge Neutral Pseudoscalar Mesons

- Isospin ($SU(2)$ -flavour) breaking is determined by the current-mass difference $m_u - m_d$, while $SU(3)$ -flavour breaking can be measured via $m_s - (m_u + m_d)/2$.



Charge Neutral Pseudoscalar Mesons

- Isospin ($SU(2)$ -flavour) breaking is determined by the current-mass difference $m_u - m_d$, while $SU(3)$ -flavour breaking can be measured via $m_s - (m_u + m_d)/2$.
- Determine effect of these differences in current-quark mass throughout the hadron spectrum, which leads one to consider the difference in mass between charged and neutral hadrons.



Charge Neutral Pseudoscalar Mesons

- Isospin ($SU(2)$ -flavour) breaking is determined by the current-mass difference $m_u - m_d$, while $SU(3)$ -flavour breaking can be measured via $m_s - (m_u + m_d)/2$.
- Determine effect of these differences in current-quark mass throughout the hadron spectrum, which leads one to consider the difference in mass between charged and neutral hadrons.
- Part of that splitting is electromagnetic, but constraining the strong component is necessary before one can know just how large that electromagnetic contribution might be.



Charge Neutral Pseudoscalar Mesons

- Isospin ($SU(2)$ -flavour) breaking is determined by the current-mass difference $m_u - m_d$, while $SU(3)$ -flavour breaking can be measured via $m_s - (m_u + m_d)/2$.
- Determine effect of these differences in current-quark mass throughout the hadron spectrum, which leads one to consider the difference in mass between charged and neutral hadrons.
- In considering neutral pseudoscalars, the $U(N_f)$ axial-vector Ward-Takahashi identity becomes relevant.
 - It necessarily includes a contribution from the non-Abelian anomaly.
 - A discussion of $\eta-\eta'$ splitting is therefore unavoidable.



Charge Neutral Pseudoscalar Mesons

$$\begin{aligned} P_\mu \Gamma_{5\mu}^a(k; P) &= \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) \\ &\quad - 2i\mathcal{M}^{ab}\Gamma_5^b(k; P) - \mathcal{A}^a(k; P) \end{aligned}$$



Charge Neutral Pseudoscalar Mesons

$$\begin{aligned} P_\mu \Gamma_{5\mu}^a(k; P) &= \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) \\ &\quad - 2i\mathcal{M}^{ab}\Gamma_5^b(k; P) - \mathcal{A}^a(k; P) \end{aligned}$$

- $\{\mathcal{F}^a | a = 0, \dots, N_f^2 - 1\}$ are the generators of $U(N_f)$



Charge Neutral Pseudoscalar Mesons

$$\begin{aligned} P_\mu \Gamma_{5\mu}^a(k; P) &= \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) \\ &\quad - 2i\mathcal{M}^{ab}\Gamma_5^b(k; P) - \mathcal{A}^a(k; P) \end{aligned}$$

- $\{\mathcal{F}^a | a = 0, \dots, N_f^2 - 1\}$ are the generators of $U(N_f)$
- $\mathcal{S} = \text{diag}[S_u, S_d, S_s, S_c, S_b, \dots]$



Charge Neutral Pseudoscalar Mesons

$$\begin{aligned} P_\mu \Gamma_{5\mu}^a(k; P) &= \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) \\ &\quad - 2i\mathcal{M}^{ab}\Gamma_5^b(k; P) - \mathcal{A}^a(k; P) \end{aligned}$$

- $\{\mathcal{F}^a | a = 0, \dots, N_f^2 - 1\}$ are the generators of $U(N_f)$
- $\mathcal{S} = \text{diag}[S_u, S_d, S_s, S_c, S_b, \dots]$
- $\mathcal{M}^{ab} = \text{tr}_F [\{\mathcal{F}^a, \mathcal{M}\} \mathcal{F}^b],$
 $\mathcal{M} = \text{diag}[m_u, m_d, m_s, m_c, m_b, \dots] = \text{matrix of current-quark bare masses}$



Charge Neutral Pseudoscalar Mesons

$$\begin{aligned} P_\mu \Gamma_{5\mu}^a(k; P) &= \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) \\ &\quad - 2i\mathcal{M}^{ab}\Gamma_5^b(k; P) - \mathcal{A}^a(k; P) \end{aligned}$$

- $\{\mathcal{F}^a | a = 0, \dots, N_f^2 - 1\}$ are the generators of $U(N_f)$
- $\mathcal{S} = \text{diag}[S_u, S_d, S_s, S_c, S_b, \dots]$
- $\mathcal{M}^{ab} = \text{tr}_F [\{\mathcal{F}^a, \mathcal{M}\}\mathcal{F}^b],$
 $\mathcal{M} = \text{diag}[m_u, m_d, m_s, m_c, m_b, \dots] = \text{matrix of current-quark bare masses}$
- The final term in the last line expresses the non-Abelian axial anomaly.



Charge Neutral Pseudoscalar Mesons

$$\begin{aligned} P_\mu \Gamma_{5\mu}^a(k; P) &= \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) \\ &\quad - 2i\mathcal{M}^{ab} \Gamma_5^b(k; P) - \mathcal{A}^a(k; P) \end{aligned}$$

- $\mathcal{A}^a(k; P) = \mathcal{S}^{-1}(k_+) \delta^{a0} \mathcal{A}_U(k; P) \mathcal{S}^{-1}(k_-)$
$$\mathcal{A}_U(k; P) = \int d^4x d^4y e^{i(k_+ \cdot x - k_- \cdot y)} N_f \langle \mathcal{F}^0 q(x) \mathcal{Q}(0) \bar{q}(y) \rangle$$



Charge Neutral Pseudoscalar Mesons

$$\begin{aligned} P_\mu \Gamma_{5\mu}^a(k; P) &= \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) \\ &\quad - 2i\mathcal{M}^{ab}\Gamma_5^b(k; P) - \mathcal{A}^a(k; P) \end{aligned}$$

- $\mathcal{A}^a(k; P) = \mathcal{S}^{-1}(k_+) \delta^{a0} \mathcal{A}_U(k; P) \mathcal{S}^{-1}(k_-)$
$$\mathcal{A}_U(k; P) = \int d^4x d^4y e^{i(k_+ \cdot x - k_- \cdot y)} N_f \langle \mathcal{F}^0 q(x) \mathcal{Q}(0) \bar{q}(y) \rangle$$
- $\mathcal{Q}(x) = i \frac{\alpha_s}{4\pi} \text{tr}_C [\epsilon_{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma}(x)] = \partial_\mu K_\mu(x)$

... The topological charge density operator.

(Trace is over colour indices & $F_{\mu\nu} = \frac{1}{2} \lambda^a F_{\mu\nu}^a$.)



Charge Neutral Pseudoscalar Mesons

$$\begin{aligned} P_\mu \Gamma_{5\mu}^a(k; P) &= \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) \\ &\quad - 2i\mathcal{M}^{ab}\Gamma_5^b(k; P) - \mathcal{A}^a(k; P) \end{aligned}$$

- $\mathcal{A}^a(k; P) = \mathcal{S}^{-1}(k_+) \delta^{a0} \mathcal{A}_U(k; P) \mathcal{S}^{-1}(k_-)$
$$\mathcal{A}_U(k; P) = \int d^4x d^4y e^{i(k_+ \cdot x - k_- \cdot y)} N_f \langle \mathcal{F}^0 q(x) \mathcal{Q}(0) \bar{q}(y) \rangle$$
- $\mathcal{Q}(x) = i \frac{\alpha_s}{4\pi} \text{tr}_C [\epsilon_{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma}(x)] = \partial_\mu K_\mu(x)$

... The topological charge density operator.

- Important that only $\mathcal{A}^{a=0}$ is nonzero.



Charge Neutral Pseudoscalar Mesons

$$P_\mu \Gamma_{5\mu}^a(k; P) = \mathcal{S}^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a \mathcal{S}^{-1}(k_-) - 2i\mathcal{M}^{ab} \Gamma_5^b(k; P) - \mathcal{A}^a(k; P)$$

- $\mathcal{A}^a(k; P) = \mathcal{S}^{-1}(k_+) \delta^{a0} \mathcal{A}_U(k; P) \mathcal{S}^{-1}(k_-)$
 $\mathcal{A}_U(k; P) = \int d^4x d^4y e^{i(k_+ \cdot x - k_- \cdot y)} N_f \langle \mathcal{F}^0 q(x) \mathcal{Q}(0) \bar{q}(y) \rangle$
- $\mathcal{Q}(x) = i \frac{\alpha_s}{4\pi} \text{tr}_C [\epsilon_{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma}(x)] = \partial_\mu K_\mu(x)$

... The topological charge density operator.

- Important that only $\mathcal{A}^{a=0}$ is nonzero.
- NB. While $\mathcal{Q}(x)$ is gauge invariant, the associated Chern-Simons current, K_μ , is not.
Means that *no physical* boson can couple to K_μ .



Charge Neutral Pseudoscalar Mesons

Flavour symmetry breaking and meson masses

Mandar S. Bhagwat, Lei Chang, Yu-Xin Liu, Craig D. Roberts
and Peter C. Tandy ... nucl-th/arXiv:0708.1118

- Many new results
 - Here, focus on $N_f = 3$
 - Will only describe those results which are exact in QCD
 - Omit numerous illustrations and calculations



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
... otherwise all pseudoscalar mesons are Goldstone Modes!



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
- Allow that $\Gamma_{5\mu}^0$ might possess a longitudinal massless bound-state pole:

$$\Gamma_{5\mu}^0(k; P) \Big|_{P^2 \approx 0} = r_A^0 \frac{P_\mu}{P^2} \Gamma_{BS}(k; P)$$

$$+ \mathcal{F}^0 \gamma_5 \left[\gamma_\mu F_R^0(k; P) + \gamma \cdot k k \cdot P G_R^0(k; P) + \sigma_{\mu\nu} k_\mu P_\nu H_R^0(k; P) + \tilde{\Gamma}_{5\mu}^0(k; P) \right]$$



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
- Allow that $\Gamma_{5\mu}^0$ might possess a longitudinal massless bound-state pole:

$$\Gamma_{5\mu}^0(k; P) \Big|_{P^2 \approx 0} = r_A^0 \frac{P_\mu}{P^2} \Gamma_{BS}(k; P)$$

$$+ \mathcal{F}^0 \gamma_5 \left[\gamma_\mu F_R^0(k; P) + \gamma \cdot k k \cdot P G_R^0(k; P) + \sigma_{\mu\nu} k_\mu P_\nu H_R^0(k; P) + \tilde{\Gamma}_{5\mu}^0(k; P) \right]$$

- $F_R^0, G_R^0, H_R^0, \Gamma_{5\mu}^0(k; P)$ are regular as $P^2 \rightarrow 0$
- $P_\mu \tilde{\Gamma}_{5\mu}^0 \sim \mathcal{O}(P^2)$
- $\Gamma_{BS}(k; P)$ is the possible bound-state's Bethe-Salpeter amplitude & r_A^0 is its residue



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
- Allow that $\Gamma_{5\mu}^0$ might possess a longitudinal massless bound-state pole:

$$\Gamma_{5\mu}^0(k; P) \Big|_{P^2 \approx 0} = r_A^0 \frac{P_\mu}{P^2} \Gamma_{BS}(k; P)$$

$$+ \mathcal{F}^0 \gamma_5 \left[\gamma_\mu F_R^0(k; P) + \gamma \cdot k k \cdot P G_R^0(k; P) + \sigma_{\mu\nu} k_\mu P_\nu H_R^0(k; P) + \tilde{\Gamma}_{5\mu}^0(k; P) \right]$$

- Bethe-Salpeter amplitude

$$\begin{aligned} \Gamma_{BS}(k; P) = & 2\mathcal{F}^0 \gamma_5 [iE_{BS}(k; P) + \gamma \cdot P F_{BS}(k; P) \\ & + \gamma \cdot k k \cdot P G_{BS}(k; P) + \sigma_{\mu\nu} k_\mu P_\nu H_{BS}(k; P)] . \end{aligned}$$



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
- Allow that $\Gamma_{5\mu}^0$ might possess a longitudinal massless bound-state pole:

$$\Gamma_{5\mu}^0(k; P) \Big|_{P^2 \approx 0} = r_A^0 \frac{P_\mu}{P^2} \Gamma_{BS}(k; P)$$

$$+ \mathcal{F}^0 \gamma_5 \left[\gamma_\mu F_R^0(k; P) + \gamma \cdot k k \cdot P G_R^0(k; P) + \sigma_{\mu\nu} k_\mu P_\nu H_R^0(k; P) + \tilde{\Gamma}_{5\mu}^0(k; P) \right]$$

- Bethe-Salpeter amplitude

$$\begin{aligned} \Gamma_{BS}(k; P) = & 2\mathcal{F}^0 \gamma_5 [iE_{BS}(k; P) + \gamma \cdot P F_{BS}(k; P) \\ & + \gamma \cdot k k \cdot P G_{BS}(k; P) + \sigma_{\mu\nu} k_\mu P_\nu H_{BS}(k; P)] . \end{aligned}$$

- Structure of anomaly contribution is similar

$$\begin{aligned} \mathcal{A}^0(k; P) = & \mathcal{F}^0 \gamma_5 [i\mathcal{E}_A(k; P) + \gamma \cdot P \mathcal{F}_A(k; P) \\ & + \gamma \cdot k k \cdot P \mathcal{G}_A(k; P) + \sigma_{\mu\nu} k_\mu P_\nu \mathcal{H}_A(k; P)] \end{aligned}$$

Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
- Leads to generalised Goldberger-Treiman relations

$$\begin{aligned} 2r_A^0 E_{BS}(k; 0) &= 2B_0(k^2) - \mathcal{E}_{\mathcal{A}}(k; 0), \\ F_R^0(k; 0) + 2r_A^0 F_{BS}(k; 0) &= A_0(k^2) - \mathcal{F}_{\mathcal{A}}(k; 0), \\ G_R^0(k; 0) + 2r_A^0 G_{BS}(k; 0) &= 2A'_0(k^2) - \mathcal{G}_{\mathcal{A}}(k; 0), \\ H_R^0(k; 0) + 2r_A^0 H_{BS}(k; 0) &= -\mathcal{H}_{\mathcal{A}}(k; 0), \end{aligned}$$

A_0, B_0 characterise gap equation's chiral limit solution.



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
- Leads to generalised Goldberger-Treiman relations

$$\begin{aligned} 2r_A^0 E_{BS}(k; 0) &= 2B_0(k^2) - \mathcal{E}_{\mathcal{A}}(k; 0), \\ F_R^0(k; 0) + 2r_A^0 F_{BS}(k; 0) &= A_0(k^2) - \mathcal{F}_{\mathcal{A}}(k; 0), \\ G_R^0(k; 0) + 2r_A^0 G_{BS}(k; 0) &= 2A'_0(k^2) - \mathcal{G}_{\mathcal{A}}(k; 0), \\ H_R^0(k; 0) + 2r_A^0 H_{BS}(k; 0) &= -\mathcal{H}_{\mathcal{A}}(k; 0), \end{aligned}$$

A_0, B_0 characterise gap equation's chiral limit solution.

- Follows that $\mathcal{E}_{\mathcal{A}}(k; 0) = 2B_0(k^2)$ is necessary and sufficient condition for absence of massless bound-state pole in $\Gamma_{5\mu}^0$.



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
- $\mathcal{E}_{\mathcal{A}}(k; 0) = 2B_0(k^2)$

Discussing the chiral limit

- $B_0(k^2) \neq 0$ if, and only if, chiral symmetry is dynamically broken.
- Hence, absence of massless η' bound-state is only assured through existence of intimate connection between DCSB and an expectation value of the topological charge density.



Charge Neutral Pseudoscalar Mesons

- Consider $\mathcal{A}^0 \not\equiv 0$
- $\mathcal{E}_{\mathcal{A}}(k; 0) = 2B_0(k^2)$

Discussing the chiral limit

- $B_0(k^2) \neq 0$ if, and only if, chiral symmetry is dynamically broken.
- Hence, absence of massless η' bound-state is only assured through existence of intimate connection between DCSB and an expectation value of the topological charge density.
- Further highlighted . . . we also proved

$$\begin{aligned}\langle \bar{q}q \rangle_{\zeta}^0 &= - \lim_{\Lambda \rightarrow \infty} Z_4(\zeta^2, \Lambda^2) \text{tr}_{\text{CD}} \int_q^{\Lambda} S^0(q, \zeta) \\ &= N_f \int d^4x \langle \bar{q}(x) i\gamma_5 q(x) Q(0) \rangle^0,\end{aligned}$$



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- Contribution from the meson's anomaly residue:

$$n_{\pi_i} = \sqrt{\frac{N_f}{2}} \nu_{\pi_i}, \quad \nu_{\pi_i} = \langle 0 | \mathcal{Q} | \pi_i \rangle$$

Connects bound-state to vacuum via topological charge operator



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- e.g., $N_f = 3$,
- $$m_{\eta'}^2 \begin{bmatrix} f_{\eta'}^3 \\ f_{\eta'}^8 \\ f_{\eta'}^0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ n_{\eta'} \end{bmatrix} + \begin{bmatrix} M_{3 \times 3} \end{bmatrix} \begin{bmatrix} \rho_{\eta'}^3 \\ \rho_{\eta'}^8 \\ \rho_{\eta'}^0 \end{bmatrix}.$$



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- e.g., $N_f = 3$, $m_{\eta'}^2 \begin{bmatrix} f_{\eta'}^3 \\ f_{\eta'}^8 \\ f_{\eta'}^0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ n_{\eta'} \end{bmatrix} + \begin{bmatrix} M_{3 \times 3} \end{bmatrix} \begin{bmatrix} \rho_{\eta'}^3 \\ \rho_{\eta'}^8 \\ \rho_{\eta'}^0 \end{bmatrix}$.

- Mass matrix

$$\begin{bmatrix} M_{3 \times 3} \end{bmatrix} = \begin{bmatrix} m_{110} & \sqrt{\frac{1}{3}} m_{1-10} & \sqrt{\frac{2}{3}} m_{1-10} \\ \sqrt{\frac{1}{3}} m_{1-10} & \frac{1}{3} m_{114} & \sqrt{\frac{2}{9}} m_{11-2} \\ \sqrt{\frac{2}{3}} m_{1-10} & \sqrt{\frac{2}{9}} m_{11-2} & \frac{2}{3} m_{111} \end{bmatrix},$$

$$m_{\alpha\beta\gamma} = \alpha m_u + \beta m_d + \gamma m_s;$$

$$\text{e.g., } m_{110} = m_u + m_d$$



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- e.g., $N_f = 3$,
- $$m_{\eta'}^2 \begin{bmatrix} f_{\eta'}^3 \\ f_{\eta'}^8 \\ f_{\eta'}^0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ n_{\eta'} \end{bmatrix} + \begin{bmatrix} M_{3 \times 3} \end{bmatrix} \begin{bmatrix} \rho_{\eta'}^3 \\ \rho_{\eta'}^8 \\ \rho_{\eta'}^0 \end{bmatrix}.$$

- residue projections

$$f_{\pi_i}^a P_\mu = Z_2 \text{tr} \int_q^\Lambda \mathcal{F}^a \gamma_5 \gamma_\mu \chi_{\pi_i}(q; P),$$

$$i\rho_{\pi_i}^a(\zeta) = Z_4 \text{tr} \int_q^\Lambda \mathcal{F}^a \gamma_5 \chi_{\pi_i}(q; P),$$

$$\chi_{\pi_i}(k; P) = \mathcal{S}(k_+) \Gamma_{\pi_i}(k; P) \mathcal{S}(k_-),$$

Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- e.g., $N_f = 3$,

$$m_{\eta'}^2 \begin{bmatrix} f_{\eta'}^3 \\ f_{\eta'}^8 \\ f_{\eta'}^0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ n_{\eta'} \end{bmatrix} + \begin{bmatrix} M_{3 \times 3} \end{bmatrix} \begin{bmatrix} \rho_{\eta'}^3 \\ \rho_{\eta'}^8 \\ \rho_{\eta'}^0 \end{bmatrix}.$$

- $U(N_f)$ limit: all current-quark masses = $m(\zeta)$



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- e.g., $N_f = 3$,

$$m_{\eta'}^2 \begin{bmatrix} f_{\eta'}^3 \\ f_{\eta'}^8 \\ f_{\eta'}^0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ n_{\eta'} \end{bmatrix} + \begin{bmatrix} M_{3 \times 3} \end{bmatrix} \begin{bmatrix} \rho_{\eta'}^3 \\ \rho_{\eta'}^8 \\ \rho_{\eta'}^0 \end{bmatrix}.$$

- $U(N_f)$ limit: all current-quark masses = $m(\zeta)$
 Complete content of above equation is statement

$$m_{\eta'}^2 f_{\eta'}^0 = n_{\eta'} + 2m(\zeta) \rho_{\eta'}^0(\zeta)$$



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- $U(N_f)$ limit: all current-quark masses = $m(\zeta)$

Complete content of above equation is statement

$$m_{\eta'}^2 f_{\eta'}^0 = n_{\eta'} + 2m(\zeta) \rho_{\eta'}^0(\zeta)$$

- It is argued that in QCD $n_{\eta'} \sim \frac{1}{\sqrt{N_c}}$

And it can be shown that $f_{\eta'}^0 \sim \sqrt{N_c} \sim \rho_{\eta'}^0(\zeta)$



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- $U(N_f)$ limit: all current-quark masses = $m(\zeta)$

Complete content of above equation is statement

$$m_{\eta'}^2 f_{\eta'}^0 = n_{\eta'} + 2m(\zeta) \rho_{\eta'}^0(\zeta)$$

- One thus obtains

$$m_{\eta'}^2 = \frac{n_{\eta'}}{f_{\eta'}^0} + 2m(\zeta) \frac{\rho_{\eta'}^0(\zeta)}{f_{\eta'}^0}$$



Charge Neutral Pseudoscalar Mesons

- Mass Formulae for neutral pseudoscalar mesons

$$m_{\pi_i}^2 f_{\pi_i}^a = 2 \mathcal{M}^{ab} \rho_{\pi_i}^b + \delta^{a0} n_{\pi_i}$$

- $U(N_f)$ limit: all current-quark masses = $m(\zeta)$

Complete content of above equation is statement

$$m_{\eta'}^2 f_{\eta'}^0 = n_{\eta'} + 2m(\zeta) \rho_{\eta'}^0(\zeta)$$

- One thus obtains

$$m_{\eta'}^2 = \frac{n_{\eta'}}{f_{\eta'}^0} + 2m(\zeta) \frac{\rho_{\eta'}^0(\zeta)}{f_{\eta'}^0}$$

- First term vanishes as $N_c \rightarrow \infty$ but second remains finite
- Subsequently taking $\hat{m} \rightarrow 0$, η' mass approaches zero in the manner characteristic of all **Goldstone modes**.

NB. One must take $N_c \rightarrow \infty$ before $\hat{m} \rightarrow 0$ because procedures do not commute



Charge Neutral Pseudoscalar Mesons

Flavour symmetry breaking and meson masses

Mandar S. Bhagwat, Lei Chang, Yu-Xin Liu, Craig D. Roberts
and Peter C. Tandy ... nucl-th/arXiv:0708.1118



Charge Neutral Pseudoscalar Mesons

Flavour symmetry breaking and meson masses

Mandar S. Bhagwat, Lei Chang, Yu-Xin Liu, Craig D. Roberts
and Peter C. Tandy ... nucl-th/arXiv:0708.1118

- Implications of mass formulae illustrated using elementary dynamical model, which includes *Ansatz* for that part of the Bethe-Salpeter kernel related to the non-Abelian anomaly
- The model involves two parameters, one of which is a mass-scale.



Charge Neutral Pseudoscalar Mesons

Flavour symmetry breaking and meson masses

Mandar S. Bhagwat, Lei Chang, Yu-Xin Liu, Craig D. Roberts
and Peter C. Tandy ... nucl-th/arXiv:0708.1118

- Implications of mass formulae illustrated using elementary dynamical model, which includes *Ansatz* for that part of the Bethe-Salpeter kernel related to the non-Abelian anomaly
- The model involves two parameters, one of which is a mass-scale.
- Employed in an analysis of pseudoscalar- and vector-meson bound-states



Charge Neutral Pseudoscalar Mesons

Flavour symmetry breaking and meson masses

Mandar S. Bhagwat, Lei Chang, Yu-Xin Liu, Craig D. Roberts
and Peter C. Tandy . . . nucl-th/arXiv:0708.1118

- Implications of mass formulae illustrated using elementary dynamical model, which includes *Ansatz* for that part of the Bethe-Salpeter kernel related to the non-Abelian anomaly
- The model involves two parameters, one of which is a mass-scale.
- Despite its simplicity, model is elucidative and phenomenologically efficacious; e.g., it predicts
 - $\eta - \eta'$ mixing angles of $\sim -15^\circ$ (Expt.: $-13.3^\circ \pm 1.0^\circ$)
 - $\pi^0 - \eta$ angles of $\sim 1.2^\circ$ (Expt. $p d \rightarrow {}^3\text{He} \pi^0$: $0.6^\circ \pm 0.3^\circ$)
 - Strong neutron-proton mass difference of 5.3 MeV (Lattice-QCD: 2.26 ± 0.72)



New Challenges



New Challenges

- Next Steps ... Applications to excited states and axial-vector mesons, e.g., will improve understanding of confinement interaction between light-quarks.




New Challenges

- Next Steps ... Applications to excited states and axial-vector mesons, e.g., will improve understanding of confinement interaction between light-quarks.
- Move on to the problem of a **symmetry preserving** treatment of hybrids and exotics.




New Challenges

- Another Direction . . . Also want/need information about three-quark systems



New Challenges

- Another Direction . . . Also want/need information about three-quark systems




- With this problem . . . current expertise at approximately same point as studies of mesons in 1995.



New Challenges

- Another Direction . . . Also want/need information about three-quark systems



- With this problem . . . current expertise at approximately same point as studies of mesons in 1995.
- Namely . . . Model-building and Phenomenology, constrained by the DSE results outlined already.






Nucleon EM Form Factors: A Précis

Höll, Kloker, et al.: nu-th/0412046 & nu-th/0501033





Nucleon EM Form Factors: A Précis

Höll, Kloker, et al.: nu-th/0412046 & nu-th/0501033

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons



Nucleon EM Form Factors: A Précis

Höll, Kloker, et al.: nu-th/0412046 & nu-th/0501033

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation



Nucleon EM Form Factors: A Précis

Höll, Kloker, et al.: nu-th/0412046 & nu-th/0501033

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
 ⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)
 Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$



Nucleon EM Form Factors: A Précis

Höll, Kloker, et al.: nu-th/0412046 & nu-th/0501033

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)
Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

(Oettel, Hellstern, Alkofer, Reinhardt: nucl-th/9805054)



Nucleon EM Form Factors: A Précis

Höll, Kloker, et al.: nu-th/0412046 & nu-th/0501033

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)
Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

- **But** is that good?



Nucleon EM Form Factors: A Précis

Höll, Kloker, et al.: nu-th/0412046 & nu-th/0501033

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
 ⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)
 Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

- But is that good?
 - Cloudy Bag: $\delta M_+^{\pi-\text{loop}} = -300$ to -400 MeV!



Nucleon EM Form Factors: A Précis

Höll, Kloker, et al.: nu-th/0412046 & nu-th/0501033

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
 ⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)
 Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$


- But is that good?
 - Cloudy Bag: $\delta M_+^{\pi\text{-loop}} = -300$ to -400 MeV!
 - Critical to anticipate pion cloud effects
- Roberts, Tandy, Thomas, et al., nu-th/02010084




Faddeev equation



Faddeev equation




Faddeev equation



- Linear, Homogeneous Matrix equation
 - Yields *wave function* (*Poincaré Covariant Faddeev Amplitude*) that describes quark-diquark relative motion within the nucleon
- Scalar and Axial-Vector Diquarks ... In Nucleon's Rest Frame *Amplitude* has ... *s-*, *p-* & *d-**wave* correlations



Diquark correlations



QUARK-QUARK

Dyson Schwinger Equations – Achievements and Challenges
Peking University, Beijing, China, 13-17 August, 2007 – p. 36/46



First

Contents

Back

Conclusion


Diquark correlations

- Same interaction that describes mesons also generates three coloured quark-quark correlations:
blue-red, blue-green,
green-red

- Confined ... Does not escape from within baryon.
- Scalar is isosinglet,
Axial-vector is isotriplet
- DSE and lattice-QCD


$$m_{[ud]_0+} = 0.74 - 0.82$$

$$m_{(uu)_1+} = m_{(ud)_1+} = m_{(dd)_1+} = 0.95 - 1.02$$



Harry Lee

Pions and Form Factors



Pions and Form Factors


- Dynamical coupled-channels model . . . Analyzed extensive JLab data . . . Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C 54, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C 63, 055201/1-13 (2001)



Pions and Form Factors

- Dynamical coupled-channels model . . . Analyzed extensive JLab data . . . Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C 54, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C 63, 055201/1-13 (2001)
- Pion cloud effects are large in the low Q^2 region.

Ratio of the M1 form factor in $\gamma N \rightarrow \Delta$ transition and proton dipole form factor G_D . Solid curve is $G_M^(Q^2)/G_D(Q^2)$ including pions; Dotted curve is $G_M(Q^2)/G_D(Q^2)$ without pions.*




Pions and Form Factors

- Dynamical coupled-channels model . . . Analyzed extensive JLab data . . . Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C 54, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C 63, 055201/1-13 (2001)
- Pion cloud effects are large in the low Q^2 region.

Ratio of the M1 form factor in $\gamma N \rightarrow \Delta$ transition and proton dipole form factor G_D . Solid curve is $G_M^(Q^2)/G_D(Q^2)$ including pions; Dotted curve is $G_M(Q^2)/G_D(Q^2)$ without pions.*

Quark Core

- Responsible for only 2/3 of result at small Q^2
- Dominant for $Q^2 > 2 - 3 \text{ GeV}^2$



Results: Nucleon and Δ Masses



Results: Nucleon and Δ Masses

Mass-scale parameters (in GeV) for the scalar and axial-vector diquark correlations, fixed by fitting nucleon and Δ masses



Set A – fit to the actual masses was required; whereas for

Set B – fitted mass was offset to allow for “ π -cloud” contributions

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.79	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$

- $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$



Results: Nucleon and Δ Masses

Mass-scale parameters (in GeV) for the scalar and axial-vector diquark correlations, fixed by fitting nucleon and Δ masses



Set A – fit to the actual masses was required; whereas for
Set B – fitted mass was offset to allow for “ π -cloud” contributions

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.79	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$

- $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$
- Axial-vector diquark provides significant attraction



Argonne
NATIONAL
LABORATORY

Results: Nucleon and Δ Masses


Mass-scale parameters (in GeV) for the scalar and axial-vector diquark correlations, fixed by fitting nucleon and Δ masses



Set A – fit to the actual masses was required; whereas for
Set B – fitted mass was offset to allow for “ π -cloud” contributions

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.79	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$

- $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$
- **Constructive Interference**: 1^{++} -diquark + $\partial_\mu \pi$



Angular Momentum Rest Frame

M. Oettel, et al.
nucl-th/9805054

Crude estimate based on magnitudes \Rightarrow probability for a u -quark to carry the proton's spin is $P_{u\uparrow} \sim 80\%$, with

$P_{u\downarrow} \sim 5\%$, $P_{d\uparrow} \sim 5\%$,

$P_{d\downarrow} \sim 10\%$.

Hence, by this reckoning $\sim 30\%$ of proton's rest-frame spin is located in dressed-quark angular momentum.

Nucleon-Photon Vertex



M. Oettel, M. Pichowsky
and L. von Smekal, nu-th/9909082

6 terms ...

Nucleon-Photon Vertex







constructed systematically ... current conserved automatically
for on-shell nucleons described by Faddeev Amplitude




6 terms ...

Nucleon-Photon Vertex

constructed systematically ... current conserved automatically
for on-shell nucleons described by Faddeev Amplitude




Form Factor Ratio: GE/GM



Form Factor Ratio: **GE/GM**


- Combine these elements ...



Form Factor Ratio: GE/GM

- Combine these elements ...


- Dressed-Quark Core



Form Factor Ratio: GE/GM

- Combine these elements ...


- Dressed-Quark Core
- Ward-Takahashi
Identity preserving
current*



Form Factor Ratio: GE/GM

- Combine these elements ...


- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution



Form Factor Ratio: GE/GM

- Combine these elements ...

- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution



Form Factor Ratio: GE/GM


- Combine these elements ...

- Dressed-Quark Core

- Ward-Takahashi*
Identity preserving
current

- Anticipate and
Estimate Pion
Cloud's Contribution

- All parameters fixed in
other applications ... **Not varied.**



Form Factor Ratio: GE/GM


- Combine these elements ...

- Dressed-Quark Core

- Ward-Takahashi*
Identity preserving
current

- Anticipate and
Estimate Pion
Cloud's Contribution

- All parameters fixed in
other applications ... **Not varied.**
- Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$




Form Factor Ratio: GE/GM

- Combine these elements ...

- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution

- All parameters fixed in
other applications ... **Not varied.**
 - Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$
 - Correlations in Faddeev amplitude – quark orbital
angular momentum – essential to that agreement




Form Factor Ratio: GE/GM


- Combine these elements ...

- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution

- All parameters fixed in other applications ... Not varied.
 - Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$
 - Correlations in Faddeev amplitude – quark orbital angular momentum – essential to that agreement
 - Predict Zero at $Q^2 \approx 6.5 \text{ GeV}^2$



Neutron Form Factors



Argonne
NATIONAL
LABORATORY

First


Contents

Back

Conclusion

Neutron Form Factors

- Expt. Madey, et al. nu-ex/0308007




Neutron Form Factors

- Expt. Madey, et al. nu-ex/0308007
- Calc. Bhagwat, et al. nu-th/0610080

$$\mu_p \frac{G_E^n(Q^2)}{G_M^n(Q^2)} = -\frac{r_n^2}{6} Q^2$$

Valid for $r_n^2 Q^2 \lesssim 1$




Neutron Form Factors

- Expt. Madey, et al. nu-ex/0308007
- Calc. Bhagwat, et al. nu-th/0610080

$$\mu_p \frac{G_E^n(Q^2)}{G_M^n(Q^2)} = -\frac{r_n^2}{6} Q^2$$

Valid for $r_n^2 Q^2 \lesssim 1$



- No sign yet of a zero in $G_E^n(Q^2)$, even though calculation predicts $G_E^n(Q^2 \approx 6.5 \text{ GeV}^2) = 0$
- Data to $Q^2 = 3.4 \text{ GeV}^2$ is being analysed (JLab E02-013)



Epilogue



First

Contents

Back

Conclusion

... tell everyone I'm
sorry about
EVERYTHING



Epilogue



First

Contents

Back

Conclusion

... tell everyone I'm
sorry about
EVERYTHING



Epilogue

- DCSB exists in QCD.



Office of
Science

U.S. DEPARTMENT OF ENERGY



Office of Nuclear Physics

Exploring Nuclear Matter - Quarks to Stars



... tell everyone I'm
sorry about
EVERYTHING



Epilogue

- DCSB exists in QCD.
 - It is manifest in the dressed light-quark propagator.
 - It impacts dramatically upon observables.



... tell everyone I'm
sorry about
EVERYTHING



Epilogue

- DCSB exists in QCD.
 - It is manifest in the dressed light-quark propagator.
 - It impacts dramatically upon observables.
- Confinement



... tell everyone I'm
sorry about
EVERYTHING



Epilogue

- DCSB exists in QCD.
 - It is manifest in the dressed light-quark propagator.
 - It impacts dramatically upon observables.
- Confinement
 - Can be realised in dressed propagators of elementary excitations
 - Observables can be used to explore model realisations



... tell everyone I'm
sorry about
EVERYTHING



Epilogue

- DCSB exists in QCD.
 - It is manifest in the dressed light-quark propagator.
 - It impacts dramatically upon observables.
- Confinement
 - Can be realised in dressed propagators of elementary excitations
 - Observables can be used to explore model realisations
- An excellent way to test conjectures and constrain the possibilities



... tell everyone I'm
sorry about
EVERYTHING



Epilogue

- DCSB exists in QCD.
 - It is manifest in the dressed light-quark propagator.
 - It impacts dramatically upon observables.
- Confinement
 - Can be realised in dressed propagators of elementary excitations
 - Observables can be used to explore model realisations
- An excellent way to test conjectures and constrain the possibilities
- Physics is an Experimental science




Contents

1. Nucleon Form Factors
2. Proton FF, Reprise
3. Modern Miracles
4. Dichotomy of the Pion
5. QCD's Emergent Phenomena
6. What's the Problem?
7. Dyson-Schwinger Equations
8. Persistent Challenge
9. Dressed-Quark Propagator
10. Quenched-QCD Dressed-Quark
11. Critical Mass
12. Quark σ -term
13. Hadrons
14. Bethe-Salpeter Kernel
15. Excitations& Chiral Symmetry
16. Radial Excitations
17. Radial Excitations& Lattice-QCD
18. Pion $J = 0$
19. Charge Neutral
20. Nucleon EM Form Factors
21. Faddeev equation
22. Diquark correlations
23. Pions and Form Factors
24. Results: Nucleon & Δ Masses
25. Angular Momentum
26. Nucleon-Photon Vertex
27. Form Factor Ratio: GE/GM
28. Neutron Form Factors
29. Parametrising diquark properties
30. Contemporary Reviews



Parametrising diquark properties



Parametrising diquark properties

- Dressed-quark . . . fixed by DSE and Meson Studies
 . . . Burden, Roberts, Thomson, Phys. Lett. **B 371**, 163 (1996)



Parametrising diquark properties

- Dressed-quark . . . fixed by DSE and Meson Studies
... Burden, Roberts, Thomson, Phys. Lett. **B 371**, 163 (1996)
- Non-pointlike scalar and pseudovector colour-antitriplet diquark correlations – described by
 - Bethe-Salpeter amplitudes . . . width for each – ω_{JP}
 - Confining propagators . . . mass for each – m_{JP}



Parametrising diquark properties

- Dressed-quark . . . fixed by DSE and Meson Studies
... Burden, Roberts, Thomson, Phys. Lett. **B 371**, 163 (1996)
- Non-pointlike scalar and pseudovector colour-antitriplet diquark correlations – described by
 - Bethe-Salpeter amplitudes . . . width for each – ω_{JP}
 - Confining propagators . . . mass for each – m_{JP}

Widths fixed by “asymptotic freedom” condition –

$$\left. \frac{d}{dK^2} \left(\frac{1}{m_{JP}^2} \mathcal{F}(K^2/\omega_{JP}^2) \right)^{-1} \right|_{K^2=0} = 1 \Rightarrow \omega_{JP}^2 = \frac{1}{2} m_{JP}^2 ,$$

Only two parameters; viz., diquark “masses”: m_{JP}



Contemporary Reviews

- Dyson-Schwinger Equations: Density, Temperature and Continuum Strong QCD
C.D. Roberts and S.M. Schmidt, nu-th/0005064,
Prog. Part. Nucl. Phys. **45** (2000) S1
- The IR behavior of QCD Green's functions: Confinement, DCSB, and hadrons . . .
R. Alkofer and L. von Smekal, he-ph/0007355,
Phys. Rept. **353** (2001) 281
- Dyson-Schwinger equations: A Tool for Hadron Physics
P. Maris and C.D. Roberts, nu-th/0301049,
Int. J. Mod. Phys. **E12** (2003) pp. 297-365
- Infrared properties of QCD from Dyson-Schwinger equations.
C. S. Fischer, he-ph/0605173,
J. Phys. **G 32** (2006) pp. R253-R291
- Nucleon electromagnetic form factors
J. Arrington, C.D. Roberts and J.M. Zanotti, nucl-th/0611050,
J. Phys. **G 34** (2007) pp. S23-S52.

